

Chronic Lymphocytic Leukemia (CLL)

Baseline Flow Cytometry

PQRI Data Collection Sheet

Patient's Name	Practice Medical Record Number (MRN)	Birth Date (mm/dd/yyyy) / /	Gender <input type="checkbox"/> Male <input type="checkbox"/> Female
National Provider Identifier (NPI)		Date of Service	

Clinical Information

Billing Information

Step 1 Is patient eligible for this measure?			Code Required on Claim Form
	Yes	No	
Patient is aged 18 years and older.	<input type="checkbox"/>	<input type="checkbox"/>	Verify date of birth on claim form.
Patient has a diagnosis of CLL.	<input type="checkbox"/>	<input type="checkbox"/>	Refer to coding specifications document for list of applicable codes.
There is a CPT E/M Service Code for this visit.	<input type="checkbox"/>	<input type="checkbox"/>	
If No is checked for any of the above, STOP. Do not report a CPT category II code.			
Step 2 Does patient meet or have an acceptable reason for not meeting the measure?			Code to be Reported on Line 24D of Paper Claim Form, if Yes (or Service Line 24 of Electronic Claim Form)
Baseline Flow Cytometry Studies ¹	Yes	No	
Performed	<input type="checkbox"/>	<input type="checkbox"/>	3170F
Not performed for one of the following reasons:			
• Medical (eg, not indicated, contraindicated, other medical reason)	<input type="checkbox"/>	<input type="checkbox"/>	3170F-1P
• Patient (eg, patient declined, economic, social, religious, other patient reason)	<input type="checkbox"/>	<input type="checkbox"/>	3170F-2P
• System (eg, resources to perform the services not available, other reason attributable to health care delivery system)	<input type="checkbox"/>	<input type="checkbox"/>	3170F-3P
Document reason here and in medical chart. _____ _____			If No is checked for all of the above, report 3170F-8P (Flow cytometry studies not performed at time of diagnosis or prior to initiating treatment, reason not otherwise specified.)

¹Baseline flow cytometry studies refer to testing that is performed at time of diagnosis or prior to initiating treatment for that diagnosis. Treatment may include anti-neoplastic therapy.