



## Binge Drinking References

American Medical Association. House of Delegates, Policy: H-30.942 Screening and Brief Interventions For Alcohol Problems.

CASA (Center on Addiction and Substance Abuse at Columbia University). *Missed Opportunity: National Survey of Primary Care Physicians and Patients*. New York, NY: Columbia University, National Center on Addiction and Substance Abuse, 2000.

Conigliaro J, Delos Reyes C, Parran TV, Schulz JE. Principles of screening and early intervention. In AW Graham, TK Schultz, MF Mayo-Smith, RK Ries & BB Wilford, eds. *Principles of Addiction Medicine, Third Edition* (pp. 325-336). Chevy Chase, MD: American Society of Addiction Medicine, 2003.

Fleming MF, Mundt MP, French MT, Manwell LB, Stauffacher EA, Barry KL. Benefit-cost analysis of brief physician advice with problem drinkers in primary care settings. *Medical Care*. 2000;38:4-18.

Fleming MF, Mundt MP, French MT, Manwell LB, Stauffacher EA, Barry KL. Brief physician advice for problem drinkers: long-term efficacy and benefit-cost analysis. *Alc Clin & Exp Res*. 2002;26:36-43.

Friedmann PD, McCullough D, Chin MH, Saitz R. Screening and intervention for alcohol problems. A national survey of primary care physicians and psychiatrists. *J Gen Intern Med*. 2000;15:84-91.

Friedmann PD, Saitz R, Gogineni A, Zhang JX, Stein MD. Validation of the screening strategy in the NIAAA "Physicians' Guide to Helping Patients with Alcohol Problems". *J Stud Alcohol*. 2001;62:234-238.

Mertens JR, Weisner C, Ray GT, Fireman B, Walsh K. Hazardous drinkers and drug users in HMO primary care: prevalence, medical conditions, and costs. *Alc Clin Exp Res*. 2005;29:989-998.

Miller PM, Thomas SE, Mallin R. Patient attitudes towards self-report and biomarker alcohol screening by primary care physicians. *Alcohol and Alcoholism*. 2006;41:306-310

Naimi TS, Brewer RD, Mokdad A, Denny C, Serdula MK, Marks JS. Binge drinking among US adults. *JAMA*. 2003;289:70-75.

National Institute on Alcohol Abuse and Alcoholism. *Brief Interventions (Alcohol Alert No.66)*. Rockville, MD: National Institute on Alcohol Abuse and Alcoholism, National Institutes of Health, July 2005.



National Institute on Alcohol Abuse and Alcoholism (NIAAA). *Helping Patients Who Drink Too Much. A Clinician's Guide*. Rockville, MD: National Institute on Alcohol Abuse and Alcoholism, National Institutes of Health, 2005.

Okoro CA, Brewer RD, Naimi TS, Moriarty DG, Giles WH, Mokdad AH. Binge drinking and health-related quality of life: do popular perceptions match reality? *Am J Prev Med*. 2004;26:230-233.

Poikolainen K. Effectiveness of brief interventions to reduce alcohol intake in primary health care populations: a meta- analysis. *Prev Med* 1999;28:503-509.

Saitz R. Unhealthy alcohol use. *New England Journal of Medicine*. 2005;352:596-607.

Saitz R, Mulvey KP, Plough A, Samet JH. Physician unawareness of serious substance abuse. *Am J Drug Alcohol Abuse*. 1997;23(3):343-354.

Town M, Naimi TS, Mokdad AH, Brewer RD. Health care access among U.S. adults who drink alcohol excessively: missed opportunities for prevention. *Prev Chronic Dis*. 2006 Apr;3:A53. Epub 2006 Mar 15.

Williams EC, Kivlahan DR, Saitz R, Merrill et al. Readiness to change in primary care patients who screened positive for alcohol misuse. *Ann Fam Med*. 2006;4:213-220.