

# How to diagnose prediabetes

APR 25, 2016

**Amy Farouk**

Past News Editor

---

Studies have shown that up to 30 percent of people with prediabetes will develop type 2 diabetes within five years. At a time when one in three U.S. adults has prediabetes, it's important to identify which of your patients have this condition to help them get the interventions they need right away. Learn the ways to identify patients with prediabetes in your practice.

## Two screening options, three tests

Prediabetes is a condition in which blood glucose or hemoglobin A1C (HbA1C) levels are higher than normal but not high enough to be classified as type 2 diabetes. There are two different approaches you can take to determine when patients have prediabetes: Identification at the point of care or via an electronic health record (EHR) query that results in a listing or registry.

For point-of-care identification, use a simple algorithm to walk through the steps. The process starts with giving patients a diabetes risk assessment. If the patient is at risk and has a body mass index (BMI) of  $\geq 24$  kg/m<sup>2</sup> ( $\geq 22$  kg/m<sup>2</sup>, if Asian\*) or a history of gestational diabetes, then you should use the results of a diagnostic test to determine whether the patient has normal blood sugar levels, prediabetes or diabetes.

There are three kinds of tests you can order: HbA1C, fasting plasma glucose or oral glucose tolerance test.

## Current Procedural Terminology (CPT®) for diabetes screening tests

CPT E/M codes for prevention-related office visits		CPT codes for office-based laboratory testing	
Preventive Visit New Patient Commercial/Medicaid	99381-99387	83036QW	Office-based Hemoglobin A1C testing
Preventive Visit Established Patient Commercial/Medicaid	99391-99397	82962	Office-based finger stick glucose testing
Annual Wellness Visit Initial Medicare	G0438		
Annual Wellness Visit Subsequent Medicare	G0439		
Individual Preventive Counseling* Commercial/Medicaid	99401 – Approx 15min 99402 – Approx 30min 99403 – Approx 45min 99404 – Approx 60min		
Face-to-Face Obesity	G0447 – 15 minutes		
Counseling for Obesity† Medicare			


**These codes may be useful to report services/tests performed to screen for prediabetes and diabetes.**

\* Preventive codes 99381-99397 include counseling and cannot be combined with additional counseling codes. If significant risk factor reduction and/or behavior change counseling is provided during a problem-oriented encounter, additional preventive counseling may be billed. In this case, modifier 25 code may allow for payment for both services, although this may vary by payer. Reimbursement for this code is not guaranteed.

† Must be billed with an ICD code indicating a BMI of 30 or greater. Medicare does not allow billing for another service provided on the same day.

Diagnostic test	Normal	Prediabetes	Diabetes
HbA1C(%)	< 5.7	5.7–6.4	≥ 6.5
Fasting plasma glucose (mg/dL)	< 100	100–125	≥ 126
Oral glucose tolerance test (mg/dL)	< 140	140–199	≥ 200

image not found or type unknown



Experts recommend that you have patients complete the risk assessment before their visit and arrange for pre-visit lab testing so you can spend time talking with your patients about their results during the actual visit. (Tip: If you don't routinely employ pre-visit planning, check out the AMA's STEPS Forward™ module on pre-visit planning.)

If you choose to identify patients with prediabetes via a registry, you can do that by querying your EHR and setting up a prediabetes registry.

Patients with a BMI of  $\geq 25$  kg/m<sup>2</sup> ( $\geq 23$  kg/m<sup>2</sup> for Asians) and blood glucose or HbA1C levels in the prediabetes range qualify for an evidence-based diabetes prevention program. View an algorithm that lists the inclusion and exclusion criteria for your EHR query.

When reporting the tests and diagnoses, refer to the charts below for the appropriate codes.

## International Classification of Diseases (ICD)-9 and ICD-10 for prediabetes and diabetes screening

ICD-10 code (effective 10-01-2015)	ICD-10 code description	ICD-9 code (effective through 9-30-2015)	ICD-9 code description
Z13.1	Encounter for screening for diabetes mellitus	V77.1	Diabetes screening
R73.09	Other abnormal glucose	790.29	Abnormal glucose
R73.01	Impaired fasting glucose	790.21	Impaired fasting glucose
R73.02	Impaired glucose tolerance (oral)	790.22	Impaired glucose tolerance (oral)
R73.9	Hyperglycemia, unspecified	790.29	Other abnormal glucose NOS
E66.01	Morbid obesity due to excess calories	278.01	Morbid Obesity
E66.09	Other obesity due to excess calories	278.00	Obesity (NOS)
E66.8	Other obesity	278.00	Obesity (NOS)
E66.9	Obesity, unspecified	278.00	Obesity (NOS)
E66.3	Overweight	278.02	Overweight
Z68.3x	Body mass indexes 30.0-39.9 (adult)	V85.30-V85.39	Body mass indexes 30.0-39.9 (adult)
Z68.4x	Body mass indexes ≥40.0 (adult)	V85.41-V85.45	Body mass indexes 30.0-39.9 (adult)

## What to do following a diagnosis

An effective, evidence-based way to reduce diabetes risk is to participate in a diabetes prevention program recognized by the CDC. Such programs emphasize healthy eating and increased physical activity, and they can reduce the risk of developing diabetes by more than one-half.

The AMA and the CDC jointly offer an easy way for care teams to access practical resources: Prevent Diabetes STAT: Screen, Test, Act—Today™. This initiative can help you and your team take three simple steps to improve the health of your patients:

1. Screen patients for prediabetes using the CDC Prediabetes Screening Test or the American Diabetes Association (ADA) Diabetes Risk Test
2. Test for prediabetes using one of three blood tests
3. Act by referring patients with prediabetes to a nearby diabetes prevention program

The U.S. Department of Health and Human Services announced it soon will begin covering diabetes prevention programs for Medicare beneficiaries, making this lifestyle intervention accessible to even more people.

\*These BMI levels reflect eligibility for the National Diabetes Prevention Program, as noted in the CDC Diabetes Prevention Recognition Program Standards and Operating Procedures. The ADA encourages screening for diabetes at a BMI of  $\geq 23$  kg/m<sup>2</sup> for Asian Americans and  $\geq 25$  kg/m<sup>2</sup> for non-Asian Americans, and some programs may use the ADA screening criteria for program eligibility. Please check with your diabetes prevention program provider for their specific BMI eligibility requirements.

image not found or type unknown

